

What Research Is Being Done on Prenatal Alcohol Exposure and Fetal Alcohol Spectrum Disorders in the Russian Research Community?

Svetlana Popova^{1,2,3,4,*}, Aleksandra Yaltonskaya^{5,6}, Vladimir Yaltonsky^{5,7}, Yaroslav Kolpakov^{5,7}, Ilya Abrosimov^{5,7}, Kristina Pervakov¹, Valeria Tanner¹ and Jürgen Rehm^{1,2,4,8}

¹Social and Epidemiological Research Department, Centre for Addiction and Mental Health, 33 Russell St., Toronto, ON, Canada M5S 2S1, ²Dalla Lana School of Public Health, University of Toronto, 155 College St., Toronto, ON, Canada M5T 3M7, ³Factor-Inwentash Faculty of Social Work, University of Toronto, 246 Bloor St. W., Toronto, ON, Canada M5S 1V4, ⁴Institute of Medical Sciences, University of Toronto, 1 King's College Circle, Toronto, ON, Canada M5S 1A8, ⁵Department of Preventive Addictology, National Research Centre on Addiction, 3 Malii Mogiltsevsky by-street, Moscow 119002, Russian Federation, ⁶Department of Psychiatry, Addiction and Psychotherapy, Moscow State University of Medicine and Dentistry, 20 Delegatskaya St., Build. 1, Moscow 127473, Russian Federation, ⁷Department of Clinical Psychology, Moscow State University of Medicine and Dentistry, 20 Delegatskaya St., Build. 1, Moscow, 127473, Russian Federation and ⁸Epidemiological Research Unit, Klinische Psychologie und Psychotherapie, Technische Universität Dresden, Chemnitz St. 46, D-01187 Dresden, Germany

*Corresponding author: Tel: +1-416-535-8501; Fax: +1-416-260-4156; E-mail: lana.popova@camh.ca

(Received 30 March 2013; first review notified 24 May 2013; in revised form 11 September 2013; accepted 12 September 2013)

Abstract — **Aims:** Although Russia has one of the highest rates of alcohol consumption and alcohol-attributable burden of disease, little is known about the existing research on prenatal alcohol exposure (PAE) and Fetal Alcohol Spectrum Disorders (FASDs) in this country. The objective of this study was to locate and review published and unpublished studies related to any aspect of PAE and FASD conducted in or using study populations from Russia. **Methods:** A systematic literature search was conducted in multiple English and Russian electronic bibliographic databases. In addition, a manual search was conducted in several major libraries in Moscow. **Results:** The search revealed a small pool of existing research studies related to PAE and/or FASD in Russia (126: 22 in English and 104 in Russian). Existing epidemiological data indicate a high prevalence of PAE and FASD, which underlines the strong negative impact that alcohol has on mortality, morbidity and disability in Russia. High levels of alcohol consumption by women of childbearing age, low levels of contraception use, and low levels of knowledge by health and other professionals regarding the harmful effects of PAE put this country at great risk of further alcohol-affected pregnancies. **Conclusions:** Alcohol preventive measures in Russia warrant immediate attention. More research focused on alcohol prevention and policy is needed in order to reduce alcohol-related harm, especially in the field of FASD.

INTRODUCTION

Russia has one of the highest rates of alcohol consumption in the world. According to the *Global Burden of Disease Study (2010)*, alcohol use is the second leading risk factor contributing to the disease burden in Russia (*Institute for Health Metrics and Evaluation, 2012*). The *World Health Organization (WHO) European and Global Status Report on Alcohol and Health* reported that on average, for 2003–2005, the level of per capita consumption of pure alcohol in liters among adults older than 15 years of age reached 26.7 l (16.3 l for women and 35.4 l for men; *WHO, 2010, 2011*). In Russia, 5.8% of women and 22.1% of men are heavy episodic drinkers (i.e. consume at least 60 g or more of pure alcohol on at least one occasion weekly; *WHO, 2010, 2011*). The prevalence of alcohol use disorders is also high in Russia—2.6% for women and 16.3% for men (*WHO, 2011*). As a result, Russia is characterized as having the most risky pattern of drinking and the greatest alcohol-attributable burden of disease in the world.

A recent study conducted in Saint Petersburg and the Nizhnyi Novgorod region revealed that 89% of non-pregnant women reported consuming alcohol and 65% reported binge drinking in the past 3 months (*Balachova et al., 2012a*). Forty-seven percent of women in the Nizhny Novgorod region and 28% in Saint Petersburg reported at least one binge-drinking episode at least monthly. Women who might become pregnant consumed alcohol similar to women who were not likely to become pregnant, and 54% of women in the Nizhny Novgorod region and 32% of women in Saint Petersburg were

considered to be at risk of having an alcohol-exposed pregnancy (*Balachova et al., 2012a*).

Popovitch et al. (2004) conducted a telephone-based behavioral risk factor surveillance survey in three Russian cities in 2000–2001. A random sample of 3032 residential telephone numbers was selected and 1693 interviews were conducted among adults 25–64 years of age in each selected household: 21% of women from Moscow, 8.2% of the women in Arkhangelsk and 10.8% of women in Murmansk abstain from alcohol. From 3 to 5% of women (and about 30% of men) in the above Russian cities consume >20 g of pure alcohol per day. Among women, about 30% of consumed beverages were spirits and 40% was beer. The highest level of hazardous alcohol consumption (>20 g of pure alcohol per day) among women was reported for those between 25 and 34 years of age, the most reproductive age group (*Popovitch et al., 2004*).

An article from the mid-1990s reported that 80–94% of high-school girls consumed alcohol and in major Russian cities, girls drank at the same level or higher than boys of the same age (*Koshkina and Paronyan, 1995*).

According to the official report of the Russian National Research Center on Addiction for 2011, the prevalence of alcoholism in the general population of Russia is estimated to be 1.4% (1402 per 100,000 persons), and ~0.5% of women (505 per 100,000) have been diagnosed with alcoholism (*Koshkina et al., 2012*). A ratio of 1:5 (women:men) has been reported to exist between the genders (*Altshuler, 2010*). Given that the official figures are based on reported cases only, the actual number of cases of alcoholism among women in Russia may be higher (if unreported cases are considered; *Kirganova,*

2009). It has also been suggested that in Russia, from 1999 to 2003, the number of registered cases of alcoholism in women increased from 2.2 to 3.7% (Altshuler *et al.*, 2006).

In addition to the alarming data on alcohol consumption by women in the general population, research indicates that there is a low level of contraception use. Only 49% of all women in Russia who have ever been sexually active use contraception, and that number increases to only 51% when women in the age group of 50–54 were excluded (Barden-O’Fallon *et al.*, 2010). In another sample of 347 non-pregnant women 18–44 years of age recruited at women’s clinics, 44% of women in Saint Petersburg and 70% in the Nizhny Novgorod region were sexually active and not using contraception consistently (Balachova *et al.*, 2012a).

Due to high levels of alcohol consumption within the population, paired with low rate of contraceptive use, many Russian women of childbearing age are at risk of having an alcohol-exposed pregnancy, which may result in having a child with Fetal Alcohol Spectrum Disorder (FASD). FASD is a non-diagnostic umbrella term that is used to represent the full spectrum of birth defects that are caused by prenatal alcohol exposure (PAE), which encompasses four categorical diagnostic entities: fetal alcohol syndrome (FAS), partial FAS (pFAS), alcohol-related neurodevelopmental disorder (ARND) and alcohol-related birth defects (ARBD; Chudley *et al.*, 2005). PAE is a leading preventable cause of the birth defects, which can include physical, mental, behavioral and/or learning disabilities with possible lifelong complications. In North America, the lifetime cost for some cases of FASD has been estimated to be more than one million dollars (Popova *et al.*, 2011).

Despite the high risk of, and potentially high cost associated with, FASD in Russia, there are no official surveillance efforts targeting FASD and no social facilities for children and/or adults with FASD in the country. Furthermore, very little is known about the existing pool of research on PAE or FASD in Russia. Therefore, the purpose of this study was to perform a systematic literature review in order to locate published and unpublished studies related to any aspect of PAE and FASD that were conducted in Russia or used study populations from Russia.

METHODS

Systematic literature search

Participating researchers from Canada and from Russia simultaneously conducted a systematic literature search. The procedure varied due to the differences in the availability of sources of information. For detailed search strategy in both countries and results, see Supplementary material, File S1.

The search was not limited by language of publication and was conducted up to and including December 2012.

RESULTS

In total, 22 studies in English and 104 studies in Russian were included in the present review. The results of the search strategy are shown in Fig. 1.

The 126 examined studies were categorized into the following six major groups, those that:

- (1) assessed the prevalence of alcohol consumption during pregnancy and the clinical complications of alcohol-exposed pregnancies;
- (2) examined the prevalence of FASD in the general population, or other specific populations (e.g. orphans);
- (3) assessed the efficacy of preventive measures, methods of diagnostics and interventions for individuals with FASD;
- (4) conducted literature reviews on FASD and developed educational materials for health professionals;
- (5) clinically examined children and adults and demonstrated the lifelong polysystemic teratogenic effects of alcohol on the fetus and
- (6) experimentally studied the teratogenic effects of alcohol on laboratory animals or human biological material.

A summary of all examined studies in chronological order is available in Supplementary material, Table S1.

Studies on the prevalence of alcohol consumption during pregnancy and the clinical complications of alcohol-exposed pregnancies

Some studies reported both the prevalence of alcohol consumption and clinical complications. Therefore, these studies have been categorized by the primary purpose of the study.

- (a) The prevalence of alcohol consumption during pregnancy

The search revealed 11 studies reporting the prevalence of alcohol consumption during pregnancy (see Supplementary material, Table S1). It was found that alcohol consumption during pregnancy in Russia varies from 3.1% (Talykova *et al.*, 2007) to 83% (Kurianova *et al.*, 2006) and heavy drinking during pregnancy varies from 0.2–1% (Radzinsky, 2002) to 9.2% (Shilko *et al.*, 2011a). In a study by Kosyh *et al.* (2010) it was reported that 50% of females consumed seven or more alcoholic drinks during the 3 months prior to pregnancy. Chambers *et al.* (2006) conducted a longitudinal pregnancy outcome study in the Moscow Region of Russia and revealed that 52% used alcohol in their most recent month of pregnancy. Further, these authors reported that, of those ‘ever drinkers’, 4.8% had at least one episode of five or more standard drinks, and 10.5% had at least one episode of three or four standard drinks in the most recent month of pregnancy. The survey conducted by Kristjanson *et al.* (2007) in Saint Petersburg among 899 pregnant women revealed that 96% of them drank alcohol in the year before they became pregnant; of these women, 60% reported drinking when they knew they were pregnant, and 35% had reported drinking within the past 30 days. Among pregnant women who drank within the past 30 days, 7.4% reported having 5 or more drinks on at least 1 occasion. One recent study reported that 16% of pregnant women with disorders of the pancreas and liver practiced heavy drinking (Egorova *et al.*, 2012). The drink of choice among pregnant Russian women was reported most commonly as wine, sparkling wine and beer (Kurianova




Fig. 1. Schematic diagram of the systematic search strategy. *Articles published in iteration are excluded from the total reported number of included studies.

et al., 2006; Kurianova, 2006; Sashchenko, 2007; Kalinina *et al.*, 2012).

Several studies revealed that Russian women substantially reduce their alcohol consumption after pregnancy recognition, when compared with pre-pregnancy levels (Balachova *et al.*, 2012a; Gaidukov *et al.*, 2008); however, about 20% continued to consume alcohol after pregnancy recognition (Balachova *et al.*, 2012a). Binge drinking after pregnancy recognition declined by a factor of 10, but did not disappear completely; ~6% of women had at least one binge episode.

It is important to note that the levels of alcohol consumption during pregnancy within all of the reviewed studies were based on maternal self-reports, which are likely to be imprecise and most likely underreported due to bias (e.g. recall and social desirability biases). Furthermore, the term 'standard drink' is not a term commonly used in Russia; therefore, the respondents may have had difficulty understanding this concept. In addition, there were inconsistencies between studies in the definition of a 'binge drinking'. For example, Balachova *et al.* (2012a) defined binge drinking as 'four or more standard drinks on a single occasion', while Chambers *et al.* (2006) defined binge drinking as an episode of 'five or more drinks'.

It should also be noted that Russia is geographically large and culturally diverse; therefore, regional variations are important. Thus, findings on the prevalence and patterns of alcohol consumption during pregnancy from individual studies may not be representative of all Russian women.

(a) The clinical complications of alcohol-exposed pregnancies

There were seven studies describing the clinical complications of alcohol-exposed pregnancies in Russia (Supplementary material, Table S1). The research studies included in this analysis have reported that alcohol consumption during pregnancy increases the risk of miscarriage (Sashchenko, 2007; Komarova, 2008), fetoplacental insufficiency (Orazmuradov *et al.*, 2007), premature or accelerated delivery (Komarova, 2008; Alekseeva, 2011a,b), pathology of amniotic fluid (Komarova, 2008; Alekseeva, 2011a,b) and disturbances in the mother's immune system (Kurianova *et al.*, 2006; Sashchenko, 2007; Komarova, 2008). Several other research studies reported that alcohol consumption during pregnancy decreases the levels of zinc and copper in the blood stream of pregnant women, as well as the levels of cobalt, iodine, magnesium and phosphorus in pregnant women's hair (Ogotoeva and Borisova, 2008; Ogotoeva *et al.*, 2009a,b; Alekseeva, 2011a,b). Furthermore, these studies reported that newborns of these mothers are deficient in certain essential minerals, such as cobalt, iron, iodine and zinc, and have increased levels of lithium and nickel. Shilko *et al.* (2010c, 2011c) stated that increased levels of transforming growth factor (TGF- β 1), found in the blood of alcohol misusing pregnant women, might explain the growth retardation in newborns.

Due to the small sample sizes of the studies presented above (Orazmuradov *et al.*, 2007; Ogotoeva and Borisova, 2008; Ogotoeva *et al.*, 2009a,b; Uliyanovskaia and Solovev, 2010; Shilko *et al.*, 2010c, 2011c; Kalinina *et al.*, 2012), and the fact that the level of alcohol consumption during pregnancy was not defined, was based on self-reported data obtained via unstructured interviews, and the timing of alcohol use was not specified (i.e., first, second or third trimester; Shilko *et al.*, 2010c, 2011c; Alekseeva, 2011a,b; Alekseeva and Ivanova, 2011; Kalinina *et al.*, 2012), these results should be interpreted with caution.

Studies reporting on prevalence of FASD

In total, 17 studies reported the prevalence of FASD in different populations of Russia (see, for example, Warren *et al.*, 2001; Riley *et al.*, 2003; Grigovich *et al.*, 2006; Miller *et al.*, 2007; Malakhova *et al.*, 2008; Konovalova *et al.*, 2009) (Supplementary material, Table S1). The prevalence of FAS in the general population was estimated in one study to be 18–19 per 1000 live births (Malakhova, 2012). However, a study of 326 kindergarten-aged children of the same area did not report any cases of FAS (Bubnov, 2009, 2010). The prevalence of FAS in maternity hospitals in Saint Petersburg was reported from 0.79 to 3.62 per 1000 live births; however, in specialized neonatal clinics it was almost 10 times higher—ranging from 3 to 35 per 1000 during 2000–2009 (Palchik *et al.*, 2006; Palchik and Legonkova, 2011).

The prevalence of FAS in orphanages ranged from 46 (Legonkova, 2011; Palchik and Legonkova, 2011) to 340 (Sofronova and Palchik, 2012) per 1000 live births by Russian researchers and from 55 (Stromland *et al.*, 2005) to 330 (The St. Petersburg-USA Orphanage Research Team, 2005) per 1000 live births by international researchers. For comparison purposes, it is believed that the prevalence rate of FAS in the North American general population is ~1 per 1000 live births (Roberts and Nanson, 2000; May and Gossage, 2001).

Studies by Marincheva *et al.* (2003) reported the prevalence of FAS to be 129 per 1000 in a boarding orphanage, 58 per 1000 in boarding schools, 49 per 1000 in regular orphanages and 164 per 1000 in a school of the social welfare system. The highest prevalence of FAS was found in ‘psycho-correctional’ orphanages for children with special needs, ranging from about 427–680 per 1000 (Legonkova, 2011; Palchik and Legonkova, 2011).

The prevalence of FAS among adopted children from Russia (or in some cases, from Eastern Europe, including Russia) currently living in the USA was estimated to range from 15 to 70 per 1000 live births (Aronson, 2003; Robert *et al.*, 2009). Farina *et al.* (2004) reported that 34% of children adopted from Russia were diagnosed with ARND. However, the records of alcohol-exposed pregnancy were significantly higher and constituted 19% (Albers *et al.*, 1997) to 41% (McGuinness *et al.*, 2000) of the cases. Albers *et al.* (1997) stressed that the majority of prospective adopting parents are concerned about the high risk of FASD among children adopted from Russia, due to the widespread availability of alcohol and the limited public awareness of alcohol’s detrimental effects on the fetus.

The existing FASD prevalence estimates in Russia vary greatly from population to population and study to study. These variations may reflect not only differences in maternal

drinking behavior, but also different diagnostic criteria, methods of case ascertainment (i.e. surveillance method) and populations surveyed in the different studies.

The existing studies have a number of limitations due to (a) inadequate and/or unavailable diagnostic capacity across the country and (b) lack of nation-wide diagnostic guidelines and definitions. Further, the studies presented above have weak methodologies, small sample sizes that are restricted to sub-populations, used convenience sampling and had a low response rate. There are also inconsistencies in the use of the terms ‘incidence’ and ‘prevalence’ across the studies.

Studies on FASD prevention, diagnostics and interventions

There were five studies found on selective prevention measurement, two on interventions, and three on diagnostic procedures (Jones *et al.*, 2006; Shilko *et al.*, 2008a, 2009c, 2010b, 2011a; Kuznetsova *et al.*, 2011). (Supplementary material, Table S1).

The studies by Balachova *et al.* (2007) and Isurina *et al.* (2009) demonstrated that, in Russia, with the exception of pediatricians, professionals such as physicians, psychologists, nurses and social workers have insufficient knowledge about FASD and usually do not discuss the negative effects of alcohol consumption during pregnancy with their female patients. These studies also demonstrated that gynecologists and pediatricians often had misinformed or inconsistent attitudes toward alcohol exposure during pregnancy such as ‘it is tolerable to consume low doses of alcohol in cases involving a healthy pregnancy’ (a misinformed attitude). Pediatricians, who were characterized as having inconsistent attitudes, assumed that it is important to convince a woman to abstain from alcohol consumption during pregnancy; however, good quality alcoholic beverages can be consumed during the late stage of pregnancy. Only medical doctors specializing in addiction treatment stated that complete abstinence from alcohol during pregnancy is necessary. Despite the small number of participants, these results (Balachova *et al.*, 2007; Isurina *et al.*, 2009) indicate a lack of knowledge regarding the detrimental effects of alcohol use during pregnancy, even among highly educated individuals in Russia.

In a brief PAE prevention intervention based on evidence-based interventions using techniques of motivational interviewing (Balachova *et al.*, 2010a,b), women were counseled to choose safe contraception or complete abstinence from alcohol consumption. It was recommended that gynecologists of the maternity welfare centers conduct this intervention because Russian women considered them as a reliable source of information (Moskalenko 2002, 2008a,b; Balachova *et al.*, 2010a,b).

Balachova *et al.* (2012b) also evaluated different types of informational leaflets with positive (positive visual images, list of positive outcomes for baby’s health in case of abstinence from alcohol), negative (negative visual images, list of negative consequences of alcohol consumption during the pregnancy) and neutral information (general information about healthy life style during pregnancy). The study found that information about FASD increases the general awareness of woman of childbearing age regarding FASD, stimulates them to consider changing their behavior toward abstinence from alcohol in case of pregnancy planning or pregnancy and decreases the level of alcohol consumption among women, in

general. It was also found that women who read the leaflets with positive information had better memories and were able to reproduce factual information about FASD. However, women who read negative information were more likely to develop a strong decision to abstain from alcohol during pregnancy (Balachova *et al.*, 2012b; Regentova, 2012).

As can be seen from the above, there are few published studies on FASD prevention, diagnosis and interventions in Russia. Studies of prevention efforts targeting high-risk women are completely absent.

Only two studies concerning the treatment of individuals with FAS were found in the Russian literature. Transcranial direct current stimulation was reported to be beneficial in normalizing the altered sleep-wake cycle in 13 children with FAS (Malakhova and Bubnov, 2011; Malakhova *et al.*, 2011). Another study reported that after implementing peptide-antiminotoxic therapy on 56 children with FAS, an improvement in psychomotor development was observed in 56–77% of cases, while a positive effect of standard therapy on psychomotor development was observed in 45% cases (Khasianova, 2010). These studies should be considered with caution due to (a) the absence of an appropriate control group, (b) the lack of a clear definition of the outcomes, (c) small sample sizes and (d) lack of appropriate statistical analyses.

Literature reviews and educational materials for health professionals

The search revealed 28 literature reviews that included information on FASD for health professionals (see, for example, Petrov-Maslakov, 1961; Lezhepekova, 1981; Bakanov, 1986, 1999; Mastiukova, 1986, 1989; Skalny and Skosyreva, 1987; Radzinsky and Kostin, 2009; Popova, 2010). (Supplementary material, Table S1). The reviews provided information on a variety of topics including: the evaluation of medical views on the FASD problem through the years, descriptions of the mechanisms of the damaging effects of alcohol on a fetus, descriptions of the clinical characteristics of children with FAS (e.g. growth retardation, birth defects/abnormalities, changes in phenotype, and neurodevelopmental and intellectual disorders) and strategies for FASD prevention (see, for example, Badalyan, 1986; Tabolin and Uryvchikov, 1986; Alipov and Korkhov, 1988; Lisitcyn and Sidorov, 1990; Akhmadeeva, 1997; Ramazanova and Semiatov, 2002; Palchik and Shabalov, 2009; Balachova *et al.*, 2012c).

An educational trial was conducted to evaluate FASD training developed for obstetricians/gynecologists and pediatricians (Balachova *et al.*, 2010a,b). This study claimed that the inclusion of a 3-h FASD education module in continued medical education led to significant changes in physicians' knowledge, attitudes and skills necessary for diagnosing FASD and conducting brief prevention interventions. Furthermore, this study implemented new technologies to disseminate developed FASD education materials and designed the first internet-based FASD educational resource in the Russian language for physicians and other health professionals (www.NetFAS.net; Balachova *et al.*, 2010a,b).

Several studies discussed the ways in which alcohol influences the development of FASD such as the direct damage of alcohol on sex cells, genitals (Koshkina *et al.*, 1998; Shilko *et al.*, 2008b), indirect autoimmune effects (Shilko *et al.*, 2008b) and metabolic disturbance (Lebedev, 1974; Bakanov,

1986; Anokhina and Moskalenko, 1987; Garmasheva and Konstantinova, 1988). The negative effect of alcohol on newborns through breastfeeding has also been reported (Bisarinina and Lisitcyna, 1987; Koshkina *et al.*, 1998). Additionally, some studies discuss the role of acetaldehyde in fetal central nervous system damage (Bakanov, 1986; Anokhina and Moskalenko, 1987; Garmasheva and Konstantinova, 1988). These studies report that an increase in acetaldehyde, ethanol's metabolite, due to insufficient maternal alcohol dehydrogenase activity, is the leading damaging factor in FAS development.

Several studies underline the dose-dependent effect of alcohol on the fetus. Some studies suggest that consumption of 60–80 g of pure alcohol per day leads to FAS (Frolova and Nikolaeva, 1987), while others report that consumption of 150 g of pure alcohol per day increases the chance of FAS development by 50% (Koshkina *et al.*, 1998).

Interestingly, in the older Russian sources from the 1980s, the detrimental effects of alcohol on the fetus were described and pregnancy termination for women with alcohol dependency was strongly recommended (Skosyreva, 1980; Kirushenkov, 1986).

As can be seen from the literature reviews listed above, the majority of them are outdated—70% were published before 2000. In addition, most of the existing reviews only provide general information on FAS. Information pertaining to the practical aspects of diagnostics, treatment, care and prevention are only touched on very briefly. Thus, they are extremely limited in their clinical applicability.

Clinical studies examining children and adults

There were 22 clinical studies conducted on children affected by FASD (Supplementary material, Table S1). These studies provide a description of neurodevelopmental, mental, cardiovascular, gastrointestinal, metabolic and other disorders presented in children with PAE and/or FASD. (Semenov *et al.*, 1987; Donetc, 1992; Erokhova and Bozhenov, 1997; Grijbovski *et al.*, 2002, 2004; Khatchel and Popov, 2005, 2009a,b, 2011; Miller *et al.*, 2006; Ruchkin *et al.*, 2008; Palchik *et al.*, 2009, 2011; Kashirskaya, 2010; Khoroshkina and Krivtsova, 2010; Sheffer, 2012).

The majority of these studies have serious limitations, including: (a) small sample sizes (Shurygin, 1974; Usova *et al.*, 1981; Grechany, 2002; Legonkova and Palchik, 2009), (b) lack of a control group (Usova *et al.*, 1981; Gummel *et al.*, 2007a,b; Khatchel and Popov, 2011) and (c) a failure to properly confirm PAE and thus, a failure to properly establish an FASD diagnosis (Shurygin, 1974; Kunikovskaia, 1980; Kornilov *et al.*, 2005; Sokolovskaya *et al.*, 2009). Most importantly, in all of the existing clinical studies examining children and adults a standardized diagnostic procedure was not used/indicated.

Experimental studies with laboratory animals or human embryos

The search revealed 25 experimental research studies conducted on rats (see, for example, Skosyreva, 1973; Skosyreva *et al.*, 1973; Anokhina *et al.*, 1989; Kolomeitseva *et al.*, 1989; Maizelis *et al.*, 1989; Nozdracheva *et al.*, 1989; Zabludovsky *et al.*, 1989; Zhulin and Bazyan, 1989; Chebotar and Konopistceva, 1993; Omelianchik *et al.*, 1993; Kataeva *et al.*, 2004; Kurch, 2004, 2011; Sverdlova, 2008; Vyatchanina and

Skalny, 2009; Arzamasova, 2011; Pugach, 2012; Shabanov *et al.*, 2012) and 6 studies using human biological material (Kovetcky *et al.*, 1991a,b; Kuteneva *et al.*, 2005; Solonsky, 2006, 2008; Shushpanova and Solonsky, 2012) (Supplementary material, Table S1). The majority of these studies reported polysystemic pathology due to PAE that affects growth, organ development, metabolism, micro- and macro-element balance and neurocognitive functioning. Experimental studies of rats further reinforce the knowledge on the negative effects of alcohol consumption during pregnancy, such as the possibility of depression-like symptoms or memory disorders in the newborns (Kolomeitceva and Levina, 1989). Furthermore, studies using human biological material demonstrated that when a mother consumes alcohol the embryo may develop many different deviations in brain development (Kovetcky *et al.*, 1991a; Solonsky, 2006, Solonsky and Logvinov, 2008). The details of these studies can be found in the Supplementary material, Table S1.

Many experimental research studies have serious limitations; they used small sample sizes (Skalny *et al.*, 2001; Shilko *et al.*, 2009a,b, 2010a,c, 2011b,c), did not describe their sample(s) (Skalny *et al.*, 2009; Morozova and Popova, 2010; Kurch, 2011) and did not present statistical significance (for example, Babenko and Skalny, 1986; Artiukhina *et al.*, 1989).

DISCUSSION

Overall, in the last five decades (the first identified article is dated as 1961), 126 studies related to alcohol consumption during pregnancy and FASD were found that were published in Russia (or on Russian samples). This seems low considering Russia's large area and population [17.1 million km²; 143 million people as of 2012 according to the Federal State Statistics Service of Russian Federation (FSSS; FSSS, 2012)]. This is also surprising given that Russia has one of the highest rates of alcohol consumption in the world. This leaves us with the question 'why is there such a low interest in this issue in the Russian research community?'

The reviewed studies suffer multiple methodological limitations and weaknesses outlined in the results section and thus, should be viewed with appropriate caution. Despite the limitations, these studies demonstrate a high level of alcohol consumption among Russian women of childbearing age, a low level of contraception usage and an especially high proportion of pregnant women with a pattern of heavy drinking. Based on these findings, it is conceivable that the prevalence of FASD is likely to be high in the general population of Russia.

The present systematic literature review revealed that only a few epidemiological studies reporting the prevalence of FASD exist. However, the reported figures are not generalizable to the general Russian population due to the methodological limitations of the studies (e.g. conducted in small communities with small sample sizes, or conducted among special populations). Valid population-based epidemiological studies are needed to examine the prevalence of FASD in both the general population and populations likely to be at high risk of PAE in Russia. Such information is crucial for understanding the magnitude of the problem and for initiating preventative measures at the country level.

The present review also revealed that the existing research on the negative effects of PAE and FASD available to health

and other professionals is not only limited, but also outdated, as the majority of these studies have been published prior to 2000. Moreover, medical education rarely ever includes a course describing the negative consequences of alcohol consumption during pregnancy; most medical doctors are not trained to recognize the clinical features of FASD in Russia (Isurina *et al.*, 2009).

Aside from medical doctors, other health and human services professionals (for example, social workers) are well positioned to help prevent FASD and to intervene with individuals and families affected by FASD. A major setback is that these professionals are also lacking appropriate knowledge and skills regarding the consequences of PAE in Russia (Balachova *et al.*, 2007, 2010a,b). Given the exceptionally high prevalence of children with FASD in foster care, orphanages and other child welfare systems in Russia, there is a need for proper training for professionals in these systems in order to be able to recognize a child with behavioral and/or mental health problems that may be the result of PAE. It is important to make educational materials available to a variety of professionals—such as guidelines for diagnosing and working with individuals already affected, as well as their parents and caregivers.

As a result of the limited research and medical and other educational information, the awareness and knowledge of Russian medical professionals, psychologists, social workers and other professionals about harmful effects of alcohol consumption during pregnancy and FASD is extremely limited (Balachova *et al.*, 2007, 2010a,b; Isurina *et al.*, 2009). There is a large discrepancy between the number of individuals who need to be diagnosed with FASD and receive proper care and the number of available professionals able to provide a proper diagnosis, treatment and other support services.

There is an urgent need to reduce the harmful use of alcohol among women of childbearing age in Russia and initiate prevention strategies targeting pregnant women. Prevention efforts need to be widely spread and should target women of childbearing age and clearly state that there is no safe time to consume alcohol during pregnancy, nor is there a safe amount or type of alcohol. The present literature review presents some effective strategies that could be implemented in these types of FASD prevention programs. For instance, Moskalenko (2002, 2008a,b) advises that the main method of preventing alcohol intake during pregnancy should involve informing women through primary care physicians, and to use positive reinforcement rather than threats. Also, several studies point out that there are misconceptions about FASD, as well as lack of awareness of FASD symptomatology (Balachova *et al.*, 2007). Improving awareness could be an effective way to minimize drinking in women of childbearing age. A randomized educational trial indicated that including FASD information in standard continuing medical education courses significantly improves physicians' knowledge, attitudes and skills (Balachova *et al.*, 2010a). Another study suggests that simple interventions such as distributing leaflets containing both positive and negative information about the consequences of drinking during pregnancy increase the general awareness of women of childbearing age and their attitudes about FASD (Balachova *et al.*, 2012b; Regentova, 2012). Thus, such research studies are a valuable tool for structuring FASD prevention programs that target women who may become pregnant.

As the literature indicates, the population of alcohol-dependent women in Russia is large; however, the principles

of prevention measures and interventions for alcohol-dependent women are underrepresented in the Russian professional literature. This means that the at-risk women are unlikely to receive an intervention for FASD prophylaxis. System-wide screening strategies and targeted interventions for women of childbearing age with alcohol dependence need to be developed. These recommendations should be primary prevention priorities for substance abuse prevention programs.

To conclude, the existing PAE and FASD research underlines the strong negative impact that alcohol has on mortality, morbidity and disability in Russia (Institute for Health Metrics and Evaluation, 2012; Lim et al., 2012; Neufeld and Rehm, 2013). The regulations introduced within the last decade in Russia seem to show some positive effects on both drinking behavior and health outcomes (Neufeld and Rehm, 2013). However, there is an urgent need for further alcohol-control strategies to reduce alcohol-related harm, especially in the field of PAE and FASD.

SUPPLEMENTARY MATERIAL

Supplementary material is available at *Alcohol and Alcoholism* online.

Acknowledgements — The Canadian team acknowledges that support to the Centre for Addiction and Mental Health for the salaries of scientists and infrastructure has been provided by the Ontario Ministry of Health and Long-Term Care in Canada. The views expressed in this manuscript do not necessarily reflect those of the Ontario Ministry of Health and Long-Term Care. The Russian team thanks the National Research Center on Addiction and Moscow State University of Medicine and Dentistry named after A.I. Evdokimov, Russian Ministry of Health for the salaries of scientists and support on FASD-related initiative in Russia. They also thank all members of Russian International Coordination Council on Alcohol Damage on Health and Fetal Alcohol Spectrum Disorder for encouragement and support of FASD-related initiative in Russia. The views expressed in this manuscript do not necessarily reflect those of the Russian Ministry of Health. We also thank Dr Tatiana Balachova, Associate Professor, University of Oklahoma Health Sciences Center, USA for her helpful comments on the draft of this paper.

Funding — Funding to pay the Open Access publication charges for this article was provided by Social and Epidemiological Research Department, Centre for Addiction and Mental Health.

Conflict of interest statement. None declared.

REFERENCES

Akhmadeeva EN. (1997) Alkogolnyi sindrom ploda: obzor [Fetal alcohol syndrome: a review]. *Zdravookhranenie Bashkortostana [Healthcare of Bashkortostan]* 6:46–51.

Albers LH, Johnson DE, Hostetter MK et al. (1997) Health of children adopted from the former Soviet Union and Eastern Europe. *JAMA* 278:922–4.

Alekseeva SN. (2011a) Osobennosti sostoianiia zdorovia i mikroelementnogo statusa novorozhdennykh, rodivshikhsia u materei, upotrebliaushchikh alkogol (na primere g. Iakutsk) [Health condition characteristics and micronutrient status of newborns to alcohol consuming mothers (an example of the city of Yakutsk)]. Doctoral Dissertation. Yakutsk, Russia: Nord-Ost Federal University named after M.K. Ammosov.

Alekseeva SN. (2011b) Alkogolnyi sindrom ploda [Fetal Alcohol Syndrome]. V: XV Kongress pediatrov Rossii s mezhdunarodnym uchastiem ‘Aktualnye problemy pediatrii’ [In: XV Congress of Pediatricians of Russia with International Participation ‘Current Problems of Pediatrics’]. Moscow, Russia, 14–17 Feb 2011, p. 17.

Alekseeva SN, Ivanova ON. (2011) Abstinennyi sindrom u novorozhdennogo ot materei, upotrebliaushchikh alkogol vo vremia

beremennosti [Withdrawal syndrome in neonates born to mothers who were consuming alcohol during pregnancy]. V: Resp. nauchno-prakt. konf., posv. 55-letiiu detskoj nevrologicheskoi sluzhby RS(Ia) [In: Scientific-Practical Conference Dedicated to the 55th Anniversary of Children’s Neurological Services of Republic Sakha (Yakuty)]. Yakutsk, Russia 14–15 Apr. 2011, pp. 19–22.

Alipov VI, Korkhov VV. (1988) *Alkogolizm i materinstvo [Alcoholism and Motherhood]*. Leningrad: Znanie [Leningrad: Knowledge].

Altshuler VB. (2010) *Alkogolizm [Alcoholism]*. Moskva: GEOTAR Media [Moscow: GEOTAR Media].

Altshuler VB, Ivanets NN, Kravchenko SL. (2006) *Zhenskij alkogolizm: klinicheskie aspekty [Female Alcoholism: Clinical Aspects]*. Moskva: Genius [Moscow: Genius].

Anokhina IP, Moskalenko VD. (1987) Alkogolizm v zerkale genicheskoi nauki [Alcoholism in the mirror of Genetic Science]. V: Simpozium ‘Alkogolizm i nasledstvennost’—Mezhdunarodnye nauchnye sviazi (Vesti AN SSSR) [In: The Symposium ‘Alcoholism and Heredity’—International Scientific Relations (USSR Academy of Sciences)]. Moscow, USSR, pp. 103–8.

Anokhina IP, Khristoliubova NA, Stanishevskaya AV et al. (1989) Funkcionalnye narusheniia neiromediatornykh sistem u potomstva krysa, podvergnutykh dlitelnoi alkogolnoi intoksikacii. V *Posledstviia alkogolnoi intoksikacii dlia potomstva*, pod. red. Airapetiantc MG [Functional disorders of neurotransmitter systems in the offspring of rats exposed to long-term alcohol intoxication. In Ayrapetyants MG (ed). *The Consequences of Alcohol Intoxication for Posterity*]. Moskva: Nauka [Moscow: Science], 3–8.

Aronson J. (2003) Fetal alcohol syndrome in Russian orphanages. *Orphan Doctor* Available at: http://www.orphandocor.com/medical/commondiseases/fas/fas_russian.html (15 January 2013, date last accessed).

Artiukhina NI, Nozdracheva LV, Mikheeva TS. (1989) Postnatalnoe razvitiie kory golovnogo mozga antenatalno alkogolizirovannykh krysa. V *Posledstviia alkogolnoi intoksikacii dlia potomstva*, pod. red. Airapetiantc MG [Postnatal development of the cerebral cortex of prenatally alcohol-exposed rats. In Ayrapetyants MG (ed). *The Consequences of Alcohol Intoxication for Posterity*]. Moskva: Nauka [Moscow: Science], 9–12.

Arzamasova OA. (2011) Uglevod-belkovye komplekxy pecheni i syvorotki krovi prenatalno alkogolizirovannykh krysa [Carbohydrate-protein complexes of liver and blood serums of prenatally alcohol-exposed rats]. V: Vseross. nauchn.-prakt. konf. biokhimiikov i spetsialistov po laboratornoi meditsine ‘Meditsinskaia biokhimiia i klinicheskaia laboratornaia diagnostika v aspekte modernizatsii sistemy nauchnykh issledovanii’ [In: Russian Scientific-Practical Conference of Biochemistry Specialists and Specialists in Laboratory Medicine ‘Medical Biochemistry and Clinical Laboratory Diagnostics in the Perspective of Modernizing the System of Scientific Research’]. Omsk, Russia, 20–21 Sep 2011, pp. 21–8.

Babenko GA, Skalny AV. (1986) Vliianie khronicheskoi alkogolnoi intoksikacii na obmen tseinka u materi i potomstva [The influence of chronic alcohol intoxication on zinc metabolism in mother and offspring]. *Akusherstvo i Ginekologiya [Obstet Gynaecol]* 3:50–2.

Badalyan LO. (1986) Sovremennye problemy alkogolnoi embrio- i fetopatii [Modern problems of alcohol embryo- and fetopathy]. *Zhurn Nevri i Psikiatrii im S.S. Korsakova [J Neur Psychiatry named after S.S. Korsakov]* 86:1443–9.

Bakanov MI. (1986) Alkogol i metabolism rebenka [Alcohol and metabolism of a child]. *Pediatria [Pediatrics]* 10:52–4.

Bakanov MI. (1999) Vliianie alkogolia na obmen veshchestv u detei i podrostkov (obzor literatury) [The influence of alcohol on the metabolism of children and adolescents (literature review)]. *Detskii Doktor [Pediatrician]* 3:40–1.

Balachova TN, Bonner BL, Isurina GL et al. (2007) Use of focus groups in developing FAS/FASD prevention in Russia. *Subst Use Misuse* 42:881–94.

Balachova T, Bonner B, Isurina G et al. (2010a) Changing physicians’ knowledge, skills and attitudes to prevent FASD in Russia. Proceedings of the 33rd Annual Scientific Meeting of the Research Society on Alcoholism. *Alcohol Clin Exp Res* 34:210A.

Balachova TN, Isurina GL, Grandilevskaya IV et al. (2010b) Profilaktika alkogolnogo sindroma ploda i narusheniia neirozrav-

- tiia u detei: tekhnologiia razrabotki profilakticheskoi programmy [Prevention of Fetal Alcohol Syndrome and neurodevelopmental disorders in children: Technology of prevention program developing]. V: Mezhdunarodnaia konferentsiia po klinicheskoi psikhologii detstva 'Rozhdenie i Zhizn'. [In: International Conference on Clinical Psychology of Childhood 'Birth and Life']. St. Petersburg, Russia, 15–16 Oct 2010, pp. 233–8.
- Balachova T, Bonner BL, Chaffin M *et al.* (2012a) Women's alcohol consumption and the risk for alcohol-exposed pregnancies in Russia. *Addiction* **107**:109–17.
- Balachova TN, Bonner BL, Isurina GL *et al.* (2012b) Izuchenie vliianiia informatsionnykh materialov na otnoshenie zhenshchin k upotrebleniiu alkogolja vo vremia beremennosti [Investigation of effects of informational materials on women's attitudes to alcohol consumption during pregnancy]. *Vestnik Sankt-Peterburgskogo Universiteta. Sankt-Peterburg: Sankt-Peterburgskij Gosudarstvennyj Universitet imeni M. Lomonosova [Bulletin of St. Petersburg State University. St. Petersburg: St. Petersburg State University named after M. Lomonosov]* **12**:101–8.
- Balachova TN, Volkova EN, Isurina GL *et al.* (2012c) *Fetalnyi alkogolnyi sindrom [Fetal Alcohol Syndrome]*. Sankt-Peterburg: SPbGU [St. Petersburg: StPSU].
- Barden-O'Fallon J, Reynolds Z, Speizer IS. (2010) Women's health in the Russian Federation: the Russia Longitudinal Monitoring Survey 2010 [working paper WP-11–121]. Chapel Hill: MEASURE Evaluation.
- Bisiarina VP, Lisitsyna NF. (1987) *Predupredit bedu: alkogolizm i deti [To prevent a disaster: alcoholism and children]*. Omsk: Vsesoiuz. Dobrovol. Ob-vo Borby za Trezvost [Omsk: USSR's Voluntary Temperance Society].
- Bubnov AA. (2009) Tcirkadnyi ritm temperatury tela u detei s fetalnym alkogolnym sindromom [Circadian rhythm of the body temperature of children with Fetal Alcohol Syndrome]. V: Mezhdunarodnaia nauchno-prakticheskaiia konf. 'Aktualnye Problemy Pediatrii' [In: International Scientific-Practical Conference 'Current Issues in Paediatrics']. Ufa, Russia, 29 Oct 2009, pp. 168–71.
- Bubnov AA. (2010) Morfo-funktsionalnaia diagnostika posledstviu vnutritrobnogo alkogolnogo vozdeistviia u detei rannego vozrasta [Morpho-functional diagnostics of consequences of intra-uterine alcohol exposure in infants]. Doctoral Dissertation. Ekaterinburg, Russia: Ural's State Medical Academy.
- Chambers CD, Kavteldadze L, Joutchenko L *et al.* (2006) Alcohol consumption patterns among pregnant women in the Moscow region of the Russian Federation. *Alcohol* **38**:133–7.
- Chebotaev NA, Konopistseva LA. (1993) Mekhanizmy teratogenno deistviia alkogolia [Mechanisms of the teratogenic effects of alcohol]. *Morfologiya [Morphology]* **105**:19–26.
- Chudley A, Conry J, Cook J *et al.* (2005) Fetal alcohol spectrum disorder: Canadian guidelines for diagnosis. *CMAJ* **172**(5 suppl): S1–S21.
- Donet OY. (1992) Psikhicheskie narusheniia u potomstva materei, bolnykh khronicheskimi alkogolizmom: (klinich. i immunolog. aspekty) [Mental disorders in the offspring of mothers with chronic alcoholism: clinical and immunological aspects]. Doctoral Dissertation. Tomsk, Russia: Tomsk's Scientific Centre RAMS, SRIMH.
- Egorova AT, Kapitonov VF, Baraniuk NV *et al.* (2012) Sotsialno-gigienicheskaia kharakteristika i reproduktivnoe zdorove zhenshchin s zabolevaniiami pankreo-biliarnoi sistemy [Social and hygienic characteristics and reproductive health of women with diseases of the pancreatobiliary system]. *Sovremennye Issledovaniia Sotsialnykh Problem [Curr Res Soc Probl]* **6**:16.
- Erokhova ZN, Bozhenov Yu A. (1997) Osobennosti sostoianiia zdorovia detei rannego vozrasta s vnutritrobnou subklinicheskoi alkogolnoi intoksikatsiei [Characteristics of the health status of young children with subclinical intrauterine alcohol intoxication]. *Ros Vestn Perinatologii i Pediatrii [Rus Bull Perinatology Paediatrics]* **1**:70.
- Farina L, Leifer M, Chasnoff IJ. (2004) Attachment and behavioural difficulties in internationally adopted Russian children. *Adoption Fostering* **28**:38–49.
- Federal State Statistics Service of Russian Federation (FSSS). (2012) Available at <http://www.gks.ru/wps/wcm/connect/rosstat/rosstatsite/eng/figures/population/> (15 March 2013, date last accessed).
- Frolova OG, Nikolaeva EI. (1987) *Gigiena Beremnoi [Prenatal Hygiene]*. Moskva: Meditsina [Moscow: Medicine].
- Gaidukov SN, Nekrasov KV, Atlasov VO. (2008) Rasprostranennost upotrebleniia zhenshchinami alkogolia do i vo vremia beremennosti i ee sotsialno-demograficheskie determinanty [Prevalence of alcohol consumption among women before and during pregnancy and its social-demographic determinants]. *Zhurnal Akusherstva i Zhenskikh Boleznei [J Obstet Gynecol]* **LVII**:11–6.
- Garmasheva NL, Konstantinova NN. (1988) Povrezhdenie mozga v geneze alkogolnogo sindroma ploda [Brain damage in the genesis of fetal alcohol syndrome]. *Vestnik Akademii Meditsinskikh Nauk SSSR [Bull USSR Academy Med Sci]* **3**:59–64.
- Grechany SV. (2002) Psikhicheskoe razvitie detei rannego vozrasta, podvergnushchisia vnutritrobnomu vozdeistviu geroina i alkogolia. V *Akusherstvo i ginekologiya. Dostizheniia i puti razvitiia v 21 veke*. Sb. nauch. tr. pod red. prof. S.N. Gaidukova, posviashchennyi 75-letiiu kafedry akusherstva i ginekologii SPbGPMMA [Mental development of children who were exposed to heroin and alcohol in utero. In Gaigukov SN (ed). *Obstetrics and Gynaecology. Accomplishments and Ways of Development in 21 Century*]. Sankt-Peterburg: SPbGPMMA [St. Petersburg: StPSMPA], 52–6.
- Grigovich IN, Mebelova II, Nikula MZ. (2006) Alkogolnyi sindrom ploda [Fetal Alcohol Syndrome]. *Rossiiskii Vestnik Perinatologii i Pediatrii (Voprosy okhrany materinstva i detstva): Nauchno-prakticheskii retsenziruemyi zhurnal [Rus Bull Perinatology Paediatrics (Issues of Protection of Mother and Child): Peer-Reviewed Journal]* **51**:10–3.
- Grijbovski A, Bygren LO, Svartbo B. (2002) Socio-demographic determinants of poor infant outcome in North-West Russia. *Pediatr Perinat Epidemiol* **16**:255–62.
- Grijbovski A, Bygren LO, Svartbo B *et al.* (2004) Housing conditions, perceived stress, smoking, and alcohol: determinants of fetal growth in northwest Russia. *Acta Obstet Gynecol Scand* **83**:1159–66.
- Gummel KK, Brzheskiy VV, Igge Y. (2007a) Nekotorye kharakteristiki zritel'nogo analizatora u detei s fetalnym alkogolnym sindromom [Characteristics of the visual analyzer in children with Fetal Alcohol Syndrome]. *Russkii Meditsinskii Zhurnal. Prilozhenie 'Klinicheskaiia Oftalmologiya' [Rus Med J App 'Clin Ophthalmol']* **8**:25–8.
- Gummel KK, Brzheskiy VV, Igge Y. (2007b) Rezultaty vizometrii, refraktometrii i staticheskoi kompiuternoi perimetrii (RBP) u detei s fetalnym alkogolnym sindromom. V: *Sovremennye Problemy Oftalmologii*. Sb. Nauchnykh Statei [Results of optometry, refractometry, and statistical computerized compare rarebit perimetry (RBP) in children with fetal alcohol syndrome. In: Contemporary Problems of Ophthalmology: Collection of Scientific Articles]. Sankt-Peterburg: Piastr [St. Petersburg: Piastre], 5–8.
- Institute for Health Metrics and Evaluation. (2012) Global Burden of Disease Study Profile: Russia. Seattle, Washington DC, USA: Institute for Health Metrics and Evaluation: Available at: <http://www.healthmetricsandevaluation.org/sites/default/files/country-profiles/GBD%20Country%20Report%20-%20Russia.pdf>. (17 July 2013, date last accessed).
- Isurina GL, Pechenezhskaia MS, Regentova AY. (2009) Profilaktika alkogolnogo sindroma ploda: izuchenie ustanovok v otnoshenii upotrebleniia alkogolia vo vremia beremennosti [Prevention of Fetal Alcohol Syndrome: study of attitudes towards alcohol consumption during pregnancy]. V: *Nauchnii Simpozium 'Nasilie i Prenebrezhenie po Otnosheniiu k Detiam: Profilaktika, Vyiavlentie, Vmeshatelstvo' [In: Scientific Symposium 'Abuse and Neglect of Children: Prevention, Detection, Intervention']*. St. Petersburg, Russia, 20–22 Oct 2009, pp. 39–41.
- Jones KL, Robinson LK, Bakhireva LN *et al.* (2006) Accuracy of the diagnosis of physical features of fetal alcohol syndrome by pediatricians after specialized training. *Pediatrics* **118**:1734–8.
- Kalinina AG, Surkova LA, Zaborova IG *et al.* (2012) Vliianie razlichnykh alkogolnykh napitkov, upotrebliaemykh zhenshchinami reproduktivnogo vozrasta na strukturu patologii i zdorove novorozhdennykh [Influence of different alcoholic beverages, consumed by women of child-bearing age, on the type of pathology and the state of health of neonates]. *Narkologiya [Narcology]* **3**:43–51.
- Kashirskaya EI. (2010) Kliniko-biokhicheskaiia otenka i prognozirovanie sostoianiia zdorovia detei, razvivaiushchikhsia pod

- vozdествiem psikoaktivnykh veshchestv [Clinical-biochemical evaluation and prognosis of the health status of children exposed to psychoactive substances]. Doctoral Dissertation. Astrakhan, Russia: Astrakhan's State Medical Academy.
- Kataeva LN, Kataev SI, Krasnova TA. (2004) Vliianie alkogolnoi intoksikatsii belykh krysov na eritropoez ikh potomstva [The effect of alcohol intoxication in white rats on erythropoiesis of their breed]. *Vestnik Ivanovskogo Gosudarstvennogo Universiteta. Seriya 'Biologiya. Khimiya. Fizika. Matematika'* [Bull Ivanovo State University. Series 'Biology, Chemistry, Physics, Mathematics'] **3**:33–7.
- Khasianova TA. (2010) Otsenka effektivnosti kompleksnoi terapii fetalnogo alkogolnogo sindroma [Evaluation of the effectiveness of complex treatment of fetal alcohol syndrome]. *Uspekhi Sovremennogo Estestvoznaniia [Success Mod Nat Sci]* **7**:123–6.
- Khatckel SB, Popov IV. (2005) Malye anomalii razvitiia: prognosticheskie pokazatel' neblagopoluchiiu rebenka bolnoi alkogolizmom materi [Minor developmental anomalies: prognostic indicator of ill-being of a child to an alcohol-dependent mother]. *Morfologiya [Morphology]* **127**:68–71.
- Khatckel SB, Popov IV. (2009a) Alkogol - vazhneishii faktor fetalnogo programmirovaniia [Alcohol: the most important fetal programming factor]. V: 3ii Mezhdistsiplinarnii Rossiiskii Kongress 'Chelovek i Alkogol (Alkogolnye Bolezni)'. [In: 3rd Russian Interdisciplinary Congress 'Human and Alcohol (Alcohol-Related Diseases)']. St. Petersburg, Russia, 23–24 Apr 2009, pp. 88–9.
- Khatckel SB, Popov IV. (2009b) Vozrastnye aspekty fetalnogo programmirovaniia kardiovaskuliarnoi patologii pod vliianiem alkogolia [Age-related aspects of fetal programming of cardiovascular pathology under the influence of alcohol]. V: 3ii Mezhdistsiplinarnii Rossiiskii Kongress 'Chelovek i Alkogol (Alkogolnye Bolezni)'. [In: 3rd Russian Interdisciplinary Congress 'Human and Alcohol (Alcohol-Related Diseases)']. St. Petersburg, Russia, 23–24 Apr. 2009, pp. 89–90.
- Khatckel SB, Popov IV. (2011) Analiz fenotipa u vospitannikov domov rebenka [Analysis of the phenotype of orphans]. *Terra Medica: Peditriia [Terra Medica: Paediatrics]* **2**:35–40.
- Khoroshkina LA, Krivtsova LA. (2010) Vliianie nikotinovoi i alkogolnoi zavisimosti zhenshchiny na zdorove novorozhdennoho [Effect of women's nicotine and alcohol dependence on newborn's health]. *Mat' i Ditiia v Kuzbasse [Mother and Child in Kuzbass]* **4**:35–8.
- Kirganova VV. (2009) Mediko-sotsialnie posledstviia in'ektsionnogo upotrebleniia narkotikov v Rossii (metodi otsenki i preduprezhdeniia) [Medical and social outcomes of injected drug use in Russia (evaluation and prevention)]. Doctoral Dissertation. Moscow, Russia: NRCA.
- Kiriushchenkov AP. (1986) Alkogolnyi sindrom ploda [Fetal Alcohol Syndrome]. *Akusherstvo i Ginekologiya [Obstet Gynaecol]* **8**:5–7.
- Kolomeitseva IA, Levina OL. (1989) Vliianie alkogolnoi intoksikatsii v period vnutritrobnogo razvitiia na funktsii TsNS potomstva [Effects of alcohol intoxication during intrauterine development on CNS functions in the offspring]. *Akusherstvo i Ginekologiya [Obstet Gynaecol]* **1**:46–50.
- Kolomeitseva IA, Varmut R, Airapetiantc MG et al. (1989) Potomstvo pri eksperimentalnom alkogolizme i substanttsiia P. V *Posledstviia alkogolnoi intoksikatsii dlia potomstva*, pod.red. Airapetiantc M.G. [Breed in experimental alcoholism and P-substance. In Ayrapetyants MG (ed). *The Consequences of Alcohol Intoxication for Posterity*]. Moskva: Nauka [Moscow: Science], 33–6.
- Komarova EE. (2008) Sostoianie fetoplatcentarnoi sistemy pri upotreblenii psikoaktivnykh veshchestv (geroin i alkogol) [The condition of the fetoplacental system during consumption of psychoactive substances (heroin and alcohol)]. Doctoral Dissertation. Moscow, Russia: People Friendship University of Russia.
- Kononova VV, Kupriyanova TA, Marincheva GS. (2009) Fetalnyi alkogolnyi sindrom u detei shkolnogo vozrasta [Fetal Alcohol Syndrome in school children]. V: IV Mezhdunarodnii Kongress 'Molodoe Pokolenie XXI Veka: Aktualnye Problemy Sotsialno-psikhologicheskogo Zdorovia' [In: International Congress IV 'The Young Generation of the XXI Century: Actual Problems of Social and Mental Health']. Kirov, Russia, 22–24 Sep 2009, pp. 106–7.
- Kornilov AA, Seledtsov AM, Maksimenko AV et al. (2005) Alkogolizm roditelei i umstvennaia otstalost' ikh detei [Parental alcoholism and children's mental retardation]. *Mat' i Ditiia v Kuzbasse [Mother and Child in Kuzbass]* **1**:24–7.
- Koshkina EA, Paronyan ID. (1995) Dinamika upotrebleniia psikoaktivnykh veshchestv sredi podrostkov [Dynamics of psychoactive substance consumption among adolescents]. *Vaktualnie problemi deviantnogo povedeniia. Borba s sotsialnimi bolezniami* [In: *Actual Problems of Deviant Behavior. Combating Social Diseases*]. Moskva: Institut sociologii RAN [Moscow: RAS Institute of Sociology], 53–8.
- Koshkina EA, Gurtovenko VM, Paronyan ID et al. (1998) Posledstviia potrebleniia alkogolia dlia zhenshchin, podrostkov, detei i semi [The consequences of alcohol consumption for women, adolescents, children and family]. *Novosti Nauki i Tekhniki. Ser. Med. Vyp. 'Alkogolnaia Bolezn'*. *Razdel 4. Alkogolnyi Sindrom Ploda / VINITI [Scientific and Technical News. Set of Medical Volumes 'Alcohol Disease'. Chapter 'Fetal Alcohol Syndrome / RISTI]* **3**:9–22.
- Koshkina EA, Kirzhanova VV, Sidoryuk OV et al. (2012) Osnovniie pokazateli deiatelnosti narkologicheskoi sluzhbi v 2011g. [Report on the main performance indicators of narcology service in Russian Federation in 2011]. Moskva: NNTc Narkologii [Moscow: National Research Center of Addiction]. Available at: <http://www.nncn.ru/datadepot/dd51/020371.pdf> (7 July 2013, date last accessed).
- Kosyh EA, Balachova T, Bonner B et al. (2010) Alcohol consumption by pregnant women in the Nizhny Novgorod region, Russia. *Alcohol Clin Exp Res* **34**:111A.
- Kovetcky NS, Kononov GV, Orlovskaya DD et al. (1991a) Dizontogenez golovnogo mozga potomstva materei, upotreblivshikh alkogol v period beremennosti [Dysontogenesis of brain development in the offspring of mothers who consumed alcohol during pregnancy]. *Zhurn Nevr i i Psikhiatr im S.S. Korsakova [J Neur Psychiatry named after S.S. Korsakov]* **10**:57–63.
- Kovetcky NS, Solonsky AV, Yarygina EG. (1991b) Formirovanie sinapsov golovnogo mozga embrionov v norme i pri alkogolizme materi [Genesis of brain synapses in embryos in normal and in cases of alcoholic mothers]. *Zhurn Nevr iPsikhiatr im S.S. Korsakova [J Neur Psychiatry named after S.S. Korsakov]* **2**:91–3.
- Kristjanson AF, Wilsnack SC, Žvartau E et al. (2007) Alcohol use in pregnant and non-pregnant Russian women. *Alcohol Clin Exp Res* **31**:299–307.
- Kunikovskaiia LS. (1980) Klinicheskaia kharakteristika oligofrenii alkogolno-embrioticheskogo gena [Clinical characteristics of alcohol-embryonic origin oligophrenia]. *Zhurn Nevr i Psikhiatr im S.S. Korsakova [J Neur Psychiatry named after S.S. Korsakov]* **80**:417–22.
- Kurch NM. (2004) Morfofunktsionalnye osobennosti razvitiia podzheludochnoi zhelezy u krysov, alkogolizirovannykh v prenatalno periode [Special morphofunctional characteristics of pancreas development in prenatally alcohol exposed rats]. Doctoral Dissertation. Tomsk, Russia: Siberia's State Medical Academy.
- Kurch NM. (2011) Korrektsiia narushenii metabolizma uglevod-belkovykh kompleksov soedinitelnoi tkani podzheludochnoi zhelezy u prenatalno alkogolizirovannykh krysov [Correction of metabolic disturbances of carbohydrate-protein complexes of the pancreatic connective tissue of prenatally alcohol-exposed rats]. V: Vseross. Nauchn.-Prakt. Konf. Biokhimikov i Spetsialistov po Laboratornoi Meditsine 'Meditsinskaia biokhimiia i klinicheskaia laboratornaia diagnostika v aspekte modernizatsii sistemy nauchnykh issledovaniy' [In: Russian Scientific-Practical Conference of Biochemists and Specialists of Lab Medicine 'Medical Biochemistry and Clinical Laboratory Diagnostics in the Perspective of Modernization of System of Scientific Research']. Omsk, Russia, 20–21 Sep 2011, pp. 171–4.
- Kurianova NN. (2006) Osobennosti khronicheskoi intoksikatsii etanolom zhenshchin v usloviiakh sotsialno-ekologicheskogo stressa [Characteristics of women with chronic alcohol intoxication under conditions of social and ecological stress]. *Uspekhi Sovremennogo Estestvoznaniia [Success Mod Nat Sci]* **2**:37–8.
- Kurianova NN, Bolotnikova NI, Udodova LV. (2006) Kliniko-sotsialnaia kharakteristika zhenshchin, upotreblivshikh vo vremia beremennosti alkogol [Clinical and social characteristic

- of women consuming alcohol during pregnancy]. *Uspekhi Sovremennogo Estestvoznaniia [Success Mod Nat Sci]* 2:36.
- Kuteneva NA, Solonsky AV, Daniletc AV *et al.* (2005) Vliianie prenatalnoi alkogolizatsii na razvitiie embrionalnykh kapilliarov golovnogo mozga [Effects of prenatal alcohol exposure on the development of embryonic capillaries of the brain]. V: Biulleten sibirskoi meditsiny. Prilozhenie 1. 'Tezisy Dokladov V Sibirskogo Fiziologicheskogo Sezda' [In: Bull of Siberia's Medicine. App. 1. 'Theses of Reports of Siberia's Physiological Congress']. Tomsk, Russia, 29–30 Jun, 1 Jul 2005 3:163.
- Kuznetsova TA, Pavlikova AV, Sviridov AS. (2011) Otsenka litcevykh anomalii v diagnostike fetalnogo alkogolnogo sindroma ploda [Evaluation of facial anomalies in the diagnosis of Fetal Alcohol Syndrome]. V: Rossiiskii Forum 'Zdorove detei: profilaktika sotcialno-znachimyykh zabolevaniy. Sankt-Peterburg – 2011' [In: Russian Forum 'Children's health: prevention of socially significant diseases. St. Petersburg – 2011']. St. Petersburg, Russia, 12–13 May 2011, pp. 99–100.
- Lebedev BA. (1974) *Alkogol i semia [Alcohol and Family]*. Leningrad: Meditsina [Leningrad: Medicine].
- Lagonkova SV. (2011) Kliniko-funktsionalnaia kharakteristika fetalnogo alkogolnogo sindroma u detei rannego vozrasta [Clinical and functional characteristics of Fetal Alcohol Syndrome in early childhood]. Doctoral Dissertation. St. Peterburg, Russia: St. Peterburg's State Paediatric Medical Academy.
- Lagonkova SV, Palchik AB. (2009) Vozrastnye osobennosti manifestatsii fetalnogo alkogolnogo sindroma [Special age-related manifestations of Fetal Alcohol Syndrome]. V: 3ii Mezhdistsiplinarnii Rossiiskii Kongress 'Chelovek i Alkogol (Alkogolnye Bolezni)' [In: 3rd Russian Interdisciplinary Congress 'Human and Alcohol (Alcohol-related Diseases)']. St. Petersburg, Russia, 23–24 Apr 2009, p. 57.
- Lezhepekova LN. (1981) *Chto nado znat pri lechenii alkogolizma? [What Is Necessary to Know for Treatment of Alcoholism?]*. Leningrad: Meditsina [Leningrad: Medicine].
- Lim S, Vos T, Flaxman AD *et al.* (2012) A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380:2224–60.
- Lisitsyn YP, Sidorov PI. (1990) Meditsinskie posledstviia alkogolizma. Alkogolnyi sindrom ploda. V *Alkogolizm. Mediko-sotcialnye aspekty*, pod red. Lisitsyn YP [Medical consequences of alcohol. Fetal Alcohol Syndrome. In Lisitsyn YP (ed). *Alcoholism. Medical and Social Aspects*]. Moskva: Meditsina [Moscow: Medicine], 221–8.
- Maizelis MY, Zabludovsky AL, Shikhov NS. (1989) Eksperimentalnaia terapiia narushenii funktsii i metabolizma mozga, vyzvannykh prenatalnoi alkogolizatsiei. V *Posledstviia alkogolnoi intoksikatsii dlia potomstva*, pod red. Airapetiantc MG [Experimental therapy in the disruption of the function and metabolism of the brain induced by prenatal alcohol exposure. In Airapetyants MG (ed). *The Consequences of Alcohol Intoxication for Posterity*]. Moskva: Nauka [Moscow: Science], 42–8.
- Malakhova ZL. (2012) Kliniko-patogeneticheskie osnovy fetalnogo alkogolnogo sindroma u detei rannego vozrasta [Clinical and pathogenic basis of fetal alcohol syndrome in infants]. Doctoral Dissertation. Ekaterinburg, Russia: Ural's State Medical Academy.
- Malakhova ZL, Bubnov AA. (2011) Opyt ispolzovaniia transkraniialnoi mikropolarizatsii golovnogo mozga kak metoda abilitatsii detei rannego vozrasta s fetalnym alkogolnym sindromom [The experience of using transcranial micropolarization (TCMP) as a method of rehabilitation of infants with Fetal Alcohol Syndrome]. *Narkologiya [Narcology]* 12:61–3.
- Malakhova ZL, Bubnov AA, Efremov AV. (2008) Otsenka fenotipicheskikh i psikhologo-pedagogicheskikh kharakteristik detei v kontekste posledstviia alkogolnykh effektov ploda [Evaluation of phenotypical and psycho-educational characteristics of children in the context of the consequences of Fetal Alcohol Effects]. *Uralskii Meditsinskii Zhurnal [Ural's Med J]* 5:67–70.
- Malakhova ZL, Bubnov AA, Shukshina LA. (2011) Opyt primeneniia transkraniialnoi mikropolarizatsii golovnogo mozga u detei rannego vozrasta s fetalnym alkogolnym sindromom [The experience of applying transcranial micropolarization (TCMP) in infants with Fetal Alcohol Syndrome]. *Uralskii Meditsinskii Zhurnal [Ural's Med J]* 4:118–20.
- Marincheva GS, Riley EP, Konovalova VV *et al.* (2003) Fetalnyi alkogolnyi sindrom v razlichnykh kontingentakh detei i podrostkov [Fetal Alcohol Syndrome in different groups of children and adolescents]. *Sotcialnaia i Klinicheskaiia Psikiatriia [Soc Clin Psychiatry]* 3:17–22.
- Mastiukova EM. (1986) Rol alkogolizma roditelei v etiologii nervno-psikhicheskikh rasstroistv u detei [The role of parents' alcoholism in the etiology of psycho-neurological disorders in children]. *Defektologiya [Abnorm Psychol]* 2:77.
- Mastiukova EM. (1989) Defekty zreniia i intellekta pri alkogolnom sindrome ploda. [Defects of vision and intellect in fetal alcohol syndrome]. *Defektologiya [Abnorm Psychol]* 6:9–13.
- May PA, Gossage JP. (2001) Estimating the prevalence of fetal alcohol syndrome: a summary. *Alcohol Res Health* 25:159–67.
- McGuinness TM, McGuinness JP, Dyer JG. (2000) Risk and protective factors in children adopted from the former Soviet Union. *Pediatr Health Care* 14:109–16.
- Miller LC, Chan W, Litvinova A *et al.* (2006) Fetal alcohol spectrum disorders in children residing in Russian orphanages: a phenotypic survey. *Alcohol Clin Exp Res* 30:531–8.
- Miller LC, Chan W, Litvinova A *et al.* (2007) Medical diagnoses and growth of children residing in Russian orphanages. *Acta Paediatr* 96:1765–9.
- Morozova MV, Popova NK. (2010) Sochetannoe vliianie alkogolia i stressa v prenatalnom periode na povedenie vzroslykh myshei [The combined effects of alcohol and stress in the prenatal period on the behaviour of adult mice]. *Rossiiskii Fiziologicheskii Zhurnal im I.M. Sechenova [Rus Physiolog J named after I.M. Sechenov]* 96:1114–21.
- Moskalenko VD. (2002) Profilaktika alkogolnogo sindroma u ploda i sindroma lisheniia narkotikov u novorozhdennykh. V *Rukovodstvo po narkologii*, pod red. NN Ivantca i dr. [Prevention of fetal alcohol syndrome in fetuses and drug withdrawal syndrome in neonates. In Ivanetc NN *et al.* (eds). *Manual in Narcology*]. Moskva: ID Medpraktika [Moscow: Medpraktika], 326–8.
- Moskalenko VD. (2008a) Profilaktika alkogolnogo sindroma u ploda i sindroma lisheniia narkotikov u novorozhdennykh. V *Narkologiya: Natsionalnoe rukovodstvo*, pod red. NN Ivantca i dr. [Prevention of fetal alcohol syndrome in fetuses and drug withdrawal syndrome in neonates. In Ivanetc NN *et al.* (eds). *Narcology: National Manual*]. Moskva: GEOTAR-Media [Moscow: GEOTAR-Media], 643–5.
- Moskalenko VD. (2008b) Profilaktika alkogolnogo sindroma u ploda i sindroma lisheniia narkotikov u novorozhdennykh. V *Rukovodstvo po narkologii*, pod red. NN Ivantca i dr. [Prevention of Fetal Alcohol Syndrome in fetuses and drug withdrawal syndrome in neonates. In Ivanetc NN *et al.* (eds). *Manual in Narcology*]. Moskva: OOO Meditsinskoe Informatcionnoe Agenstvo [Moscow: Medical News Agency], 818–20.
- Neufeld M, Rehm J. (2013) Alcohol consumption and mortality in Russia since 2000: Are there any changes following the alcohol policy changes starting in 2006? *Alcohol Alcohol* 48:222–30.
- Nozdracheva LV, Artyukhina NI, Kolomeitseva IA *et al.* (1989) Korregiruiushchee deistvie dalargina na TcNS kryis pri antenatalnoi alkogolnoi intoksikatsii. V *Posledstviia alkogolnoi intoksikatsii dlia potomstva*, pod red. Airapetiantc MG [Corrective action of dalargin on the central nervous system of rats with antenatal alcohol intoxication. In Airapetyants MG (ed). *The Consequences of Alcohol Intoxication for Posterity*]. Moskva: Nauka [Moscow: Science], 59–63.
- Ogotoeva SN, Borisova NV. (2008) Mikroelementnyi sostav krovi beremennykh zhenshchin pri upotreblenii alkogolia [Micronutrient blood count in pregnant women consuming alcohol]. *Mikroelementy v Meditsine [Microelements Med]* 9:61.
- Ogotoeva SN, Barashkova NN, Borisova NV. (2009a) Vliianie 'umerennogo' upotrebleniia alkogolia vo vremia beremennosti na mikroelementnyi sostav krovi zhenshchin i sostoianie zdorovia novorozhdennykh detei [Effects of moderate alcohol consumption during pregnancy on maternal micronutrient blood count and the health status of the neonate]. *Iakutskii Meditsinskii Zhurnal [Yakuty's Med J]* 3:68–70.
- Ogotoeva SN, Borisova NV, Tumusov PS *et al.* (2009b) Osobennosti techeniia rannego neonatalnogo perioda novorozhdennykh ot materei, stradaiushchikh khronicheskim alkogolizmom [Characteristics of early neonatal development of neonates born to

- chronic alcohol-dependent mothers]. *Nauka i Obrazovanie [Sci Educ]* 2:49–51.
- Omelianchik MS, Artamonova VG, Selyavko AA. (1993) Asimetriia v okostenenii nizhnei cheliusti kak rannii simptom alkogolnogo sindroma ploda [Asymmetry in the ossification of the lower jawbone as an early symptom of Fetal Alcohol Syndrome]. *Zdravookhranenie Belarusi [Health Belarus]* 8:45–9.
- Orazmuradov AA, Radzinsky VE, Ogurtcov PP et al. (2007) Vliianie alkogolizma i tabakokurenii na techenie beremennosti i perinatalnyi iskhod rodov [The effects of alcoholism and cigarette smoking on the pregnancy and the perinatal outcome of labor]. *Narkologiya [Narcology]* 11:60–2.
- Palchik AB, Legonkova SV. (2011) Fetalnyi alkogolnyi sindrom: manifestatsiia i dinamika [Fetal alcohol syndrome: manifestation and dynamic]. *Obozrenie Psikhatrii i Med. Psikhologii im. V.M. Bekhtereva [Rev Psychiatry Med Psychology named after V.M. Bekhterev]* 3:17–20.
- Palchik AB, Shabalov NP. (2009) *Toksicheskie entsefalopatii novorozhdennykh [Toxic Encephalopathy of Newborns]*. Moskva: MEDpress-inform [Moscow: MEDpress-inform].
- Palchik AB, Fedorova LA, Legonkova SV. (2006) *Fetalnyi alkogolnyi sindrom. Metodicheskie rekomendatsii [Fetal Alcohol Syndrome. Medical Recommendations]*. Sankt-Peterburg: OOO ANTT-Print [St. Petersburg: ANTT-Print].
- Palchik AB, Nazdzhanova ZG, Legonkova SV. (2009) Osobennosti formirovaniia bioelektricheskoi aktivnosti u detei rannego vozrasta s antenatalnoi ekspozitciei toksichnykh veshchestv [Characteristics of the formation of bio-electrical activity in children with antenatal exposure to toxic substances]. V: 3ii Mezhdistsiplinarii Rossiiskii Kongress ‘Chelovek i Alkogol (Alkogolnye Bolezni)’ [In: 3rd Russian Interdisciplinary Congress ‘Human and Alcohol (Alcohol-related Diseases)’]. St. Petersburg, Russia, 23–24 Apr 2009, p. 62.
- Palchik AB, Legonkova SV, Sofronova GI. (2011) Etnicheskie osobennosti manifestatsii fetalnogo alkogolnogo sindroma [Ethnic characteristics of the manifestation of Fetal Alcohol Syndrome]. *Iakutskii Meditsinskii Zhurnal [Yakuty's Med J]* 2:22–5.
- Petrov-Maslakov MA. (1961) *Gigiiena zhenshchiny. V Istoricheskoe razvitiie akusherstva i ginekologii. Embriogenez. Anatomii i fiziologii zhenskikh polovykh organov. Gigiiena zhenshchiny*, pod red.toma prof. FA Syrovatko. V *Mnogotomnoe rukovodstvo po akusherstvu i ginekologii*, otv.red. prof. LS Persianov. [Hygiene of women. In: Syrovatko FA (ed). *Historical Development of Obstetrics and Gynecology. Embryogenesis. Anatomy and Physiology of the Female Genital Organs. Hygiene of Woman*. In: Persianov LS (ed). *Multivolume Manual on Obstetrics and Gynecology*]. Moskva: MEDGIZ [Moscow: Medical State Publisher], 416.
- Popova EN. (2010) *Ultrastruktura mozga.alkogol i potomstvo [Ultrastructure of the Brain, Alcohol and the Offspring]*. Moskva: Nauchnyi Mir [Moscow: Scientific World].
- Popova S, Stade B, Bekmuradov D et al. (2011) What do we know about the economic impact of Fetal Alcohol Spectrum Disorder? A systematic literature review. *Alcohol Alcohol* 46:490–7.
- Popovitch MV, Glazunov RA, Potemkin AD et al. (2004) Potreblenie alkogolia v Rossii [Alcohol consumption in Russia]. *Demoskop Weekly [Demoskop Weekly]* Available at: <http://www.demoscope.ru/weekly/2004/0143/analit02.php> (15 January 2013, date last accessed).
- Pugach PV. (2012) Vliianie dlitelnosti etanolovoi intoksikatsii na krys i immunnye organy ikh potomstva [The effects of time and duration of ethanol intoxication on rats and the immune organs of their breed]. Doctoral Dissertation. St. Petersburg, Russia: St. Petersburg's State Paediatric Medical Academy.
- Radzinsky VE. (2002) *Patologicheskoe akusherstvo. V Akusherstvo. Praktikum v 3-kh chastiakh* [Pathological obstetrics. In Radzinsky VE (ed). *Obstetrics. Practical Course in 3 Parts*]. Moskva: RUDN [Moscow: People Friendship University of Russia], 545.
- Radzinsky VE, Kostin IN. (2009) *Faktory perinatalnogo riska: vrednye privychki. V Akusherskii risk. Maksimum informatsii minimum opasnosti dlia materi i mladentca* [Perinatal risk factors: addictions. In *Obstetrical Risk. Maximum Information Minimum Danger for Mother and Newborn*]. Moskva: Eksmo [Moscow: Eksmo], 81–6.
- Ramazanov LM, Semiatov SM. (2002) Alkogol i ego vliianie na plod [Alcohol and its effects on the foetus]. *Vestnik Rossiiskogo Universiteta Druzhy Narodov. Seriya: Meditsina [Bull of People Friendship University of Russia]* 1:274–81.
- Regentova AY. (2012) Informirovannost kak faktor vliianiia na otnoshenie zhenshchin k upotrebleniiu alkogolia vo vremia beremennosti [Awareness as a factor of influence on women's attitudes to alcohol consumption during pregnancy]. *Vestnik IuUGU [Bull South Ural's State University]* 31:109–14.
- Riley EP, Mattson SN, Li T-K et al. (2003) Neurobehavioral consequences of prenatal alcohol exposure: An international perspective. *Alcohol Clin Exp Res* 27:362–73.
- Robert M, Carceller A, Domken V et al. (2009) Physical and neurodevelopmental evaluation of children adopted from Eastern Europe. *Can J Clin Pharmacol* 16:432–40.
- Roberts G, Nanson J. (2000) *Best Practices. Fetal Alcohol Syndrome/ Fetal Alcohol Effects and the Effects of Other Substance Use During Pregnancy*. Ottawa, ON: Canada's Drug Strategy Division, Health Canada.
- Ruchkin V, Gilliam WS, Mayes L. (2008) Developmental pathway modeling in considering behavior problems in young Russian children. *Child Psychiatry Hum Dev* 39:49–66.
- Sashchenko AI. (2007) *Fetoplatcentarnaia sistema pri alkogolizme i tabakokurenii* [Fetoplacental system in women with alcoholism and tobacco consumption]. Doctoral Dissertation. Moscow, Russia: People Friendship University of Russia.
- Semenov AS, Sërganova TI, Skalny AV et al. (1987) O vliianii alkogolizma roditelei na razvitiie potomstva [The effect of alcoholic parents on the development of the offspring]. *Pediatriia [Paediatrics]* 4:49–53.
- Shabanov PD, Lebedev AA, Bychkov ER. (2012) Vliianie vnutritrobnogo deistviia etanola na sozrevanie monoaminergicheskikh sistem v razvivaiushchemsia mozge krys [Effects of intrauterine alcohol exposure on the monoaminergic system maturation in the rat's developing brain]. *Russ Fiziol Zhurnal im I.M. Sechenova [Rus Physiol J named after I.M. Sechenov]* 98:202–11.
- Sheffer KK. (2012) *Funkttsionalnoe sostoiianie zritel'nogo analizatora u detei podrostkovogo vozrasta s fetalnym alkogolnym sindromom* [Functional conditions of the visual analyzer in adolescents with fetal alcohol syndrome]. Doctoral Dissertation. St. Petersburg, Russia: St. Petersburg's State Paediatric Medical Academy.
- Shilko VI, Malakhova ZL, Bubnov AA et al. (2008a) Fetalnyi alkogolnyi spektr narushenii sredi vospitannikov domov rebenka [Fetal Alcohol Spectrum Disorder in orphans]. *Narkologiya [Narcology]* 11:53–6.
- Shilko VI, Malakhova ZL, Bubnov AA. (2008b) Fetalnyi alkogolnyi sindrom [Fetal alcohol syndrome]. *Nizhegorodskii Meditsinskii Zhurnal [N Novgorod's Med J]* 2:94–8.
- Shilko VI, Malakhova ZL, Bubnov AA et al. (2009a) *Kliniko-eksperimentalnye sopostavleniia pri fetalnom alkogolnom sindrome* [Clinical and experimental comparisons in fetal alcohol syndrome]. *Uralskii Meditsinskii Zhurnal: Pediatriia [Ural's Med J Paediatrics]* 7:25–8.
- Shilko VI, Malakhova ZL, Bubnov AA et al. (2009b) Fetalnyi alkogolnyi sindrom: kliniko-eksperimentalnye sopostavleniia [Fetal alcohol syndrome: clinical and experimental comparisons]. *Narkologiya [Narcology]* 8:38–40.
- Shilko VI, Malakhova ZL, Shukshina ZL. (2009c) Fetalnyi alkogolnyi spektr narushenii sredi vospitannikov sotcialnykh uchrezhdenii [Fetal Alcohol Spectrum Disorder in children from foster homes]. *Kazanskii Meditsinskii Zhurnal [Kazan's Med J]* 90:647–50.
- Shilko VI, Malakhova ZL, Bubnov AA. (2010a) Porazhenie golovnogo mozga pri fetalnom alkogolnom sindrome [Brain damage in fetal alcohol syndrome]. *Bulleten Eksperimentalnoi Biologii i Meditsiny [Bull Exp Biol Med]* 150:100–3.
- Shilko VI, Malakhova ZL, Bubnov AA et al. (2010b) Fetalnyi alkogolnyi spektr narushenii sredi vospitannikov domov rebenka [Fetal Alcohol Spectrum Disorder in orphans]. *Vestnik VolgGU [Bull Volgograd's State Med Academy]* 4:108–11.
- Shilko VI, Zilber MY, Malakhova ZL. (2010c) Transformiruiushchii faktor rosta u beremennykh zhenshchin stradaiushchikh alkogolizmom, i u detei s fetalnym alkogolnym sindromom [Transforming growth factor in pregnant alcohol dependent women and in children with fetal alcohol syndrome]. *Rossiiskii Vestnik Perinatologii i Pediatrii [Rus Bull Perinatology Paediatrics]* 6:20–2.
- Shilko VI, Malakhova ZL, Bubnov AA. (2011a) *Fetalnyi alkogolnyi sindrom: kliniko-patogeneticheskaiia kharakteristika posledstviia u*

- detei rannego vozrasta [Fetal Alcohol Syndrome: Clinical and Pathogenetic Characteristics of the Consequences in Infants]. Ekaterinburg: UrGMA [Ekaterinburg, UrGMA].
- Shilko VI, Malakhova ZL, Bubnov AA. (2011b) K patogenezu fetalnogo alkogolnogo sindroma [The pathogenesis of Fetal Alcohol Syndrome]. *Vestnik Uralskoi Meditsinskoi Akademicheskoi Nauki [Bull Ural's Med Acad Sci]* 3:55–8.
- Shilko VI, Malakhova ZL, Zilber MY. (2011c) O roli transformiruiushchego faktora rosta TGF- β 1 v razvitii fetalnogo alkogolnogo sindroma [About the role of transforming growth factor TGF- β 1 in the development of Fetal Alcohol Syndrome]. *Nauchno-prakticheskiy Zhurnal 'Kliniko-laboratornyi Konsilium' [Sci Pract J 'Clin Lab Concillium']* 1:46–8.
- Shurygin GI. (1974) Ob osobennostiakh psikhicheskogo razvitiia detei ot materei, stradiavushchikh khronicheskim alkogolizmom [About the characteristics of the mental development of children of mothers with chronic alcoholism]. *Pediatrica [Paediatrics]* 11:71–3.
- Shushpanova TV, Solonsky AV. (2012) Sinaptogenez i formirovanie benzodiazepinovykh retseptov mozga cheloveka v usloviakh prenatalnoi alkogolizatsii [Synaptogenesis and Benzodrine receptor formation in human brain under prenatal alcohol exposure]. *Zhurn Nevr i Psikiatr im S.S. Korsakova [J Neur Psychiatry named after S.S. Korsakov]* 1:60–7.
- Skalny AV, Skosyreva AM. (1987) Defitsit tsinka u materi, ploda i potomstva pri zloupotreblenii alkogolem [Zinc deficit in the mother, foetus and offspring when alcohol abuse is present]. *Akusherstvo i Ginekologiya [Obstet Gynaecol]* 4:51–6.
- Skalny AV, Kampov-Polevoi AB, Voronin AE. (2001) Vliianie tsinka na aktivnost etanolokisliaiushchikh fermentov potomkov alkogolizirovannykh krysov [Effects of zinc on ethanol acidic activity in a breed of alcohol-exposed rats]. *Mikroelementy v Meditsine [Microelements Med]* 2:21–3.
- Skalny AV, Kin KL, Chambers KD *et al.* (2009) Narushenie obmena tsinka pri prenatalnom vozdeistvii alkogolia [Zinc metabolism disturbances under prenatal alcohol exposure]. V: I Ross. Natsionalnii Kongress po Narkologii s Mezhdunarodnym Uchastiem [In: 1st Rus National Congress on Narcology with International Participation]. Moscow, Russia, 24–27 Nov 2009, pp. 18–9.
- Skosyreva AM. (1973) Vliianie etilovogo spirita na razvitie embrionov stadii organogeneza [Effects of alcohol on the development of embryos on the stage of organogenesis]. *Akusherstvo i Ginekologiya [Obstet Gynaecol]* 4:15–8.
- Skosyreva AM. (1980) Deistvie etilovogo spirita v period ontogeneza [Effects of alcohol during ontogenesis]. *Akusherstvo i Ginekologiya [Obstet Gynaecol]* 12:7–9.
- Skosyreva AM, Balika YuD, Kartasheva VE. (1973) Vliianie etilovogo spirita na razvitie embriona i ploda v eksperimente [Effects of alcohol on the development of the embryo and foetus in an experimental setting]. *Akusherstvo i Ginekologiya [Obstet Gynaecol]* 4:18–20.
- Sofronova GI, Palchik AB. (2012) Features of maternal alcohol consumption and fetal alcohol syndrome in the Sakha Republic. *J Popul Ther Clin Pharmacol* 19:e405–6.
- Sokolovskaya TA, Azarko AE, Dmitrieva OV. (2009) Vliianie alkogolizma roditelei na formirovanie invalidnosti u detei [Effects of parental alcohol-exposure on developmental disabilities in children]. *Informatcionno-analiticheskiy vestnik 'Sotsialnye Aspekty Zdorovia Naseleniia' [Inf Anal J 'Soc Aspects Popul Health']* Available at: <http://vestnik.mednet.ru/content/view/110/30/> (15 January 2013, date last accessed).
- Solonsky AV. (2006) Formirovanie krovenosnykh sosudov embrionalnogo mozga cheloveka v usloviakh prenatalnoi alkogolizatsii [Formation of blood vessels in the human embryonic brain in conditions of prenatal alcohol exposure]. *Zhurn Nevr i Psikiatr im S. S. Korsakova [J Neur Psychiatry named after S.S. Korsakov]* 12:71–3.
- Solonsky AV. (2008) Morfologicheskie zakonomernosti rannikh stadii razvitiia golovnogo mozga v usloviakh prenatalnoi alkogolizatsii [Morphological regularities of early stages of the brain development under conditions of parental alcohol exposure]. Doctoral Dissertation. Tomsk, Russia: Siberia's State Medical Academy.
- Solonsky AV, Logvinov SV. (2008) Ultrastrukturnye i morfometricheskie osobennosti sinaptogeneza mozga embriona i plodov cheloveka v usloviakh prenatalnogo vozdeistviia etanola [Ultrastructural and morphological characteristics of synaptogenesis of the brain in human embryos and fetuses during parental alcohol exposure]. *Biulleten Sibirskoi Meditsiny [Bull Siberia Med]* 7:35–9.
- Stromland K, Mattson SN, Adnams CM *et al.* (2005) Fetal alcohol spectrum disorders: an international perspective. *Alcohol Clin Exp Res* 29:1121–6.
- Sverdlova AV. (2008) Vliianie prenatalnoi etanolovoi khronicheskoi intoksikatsii na razvitie potomstva krysov v rannie sroki postnatalnogo ontogeneza (po pokazateliam nefrogeneza) [Effects of prenatal chronic alcohol intoxication on the development of offspring of rats in early stage of postnatal ontogenesis (based on the indicators of nephrogenesis)]. *Fundamentalnye Issledovaniia [Fundam Res]* 1:148–50.
- Tabolin VA, Uryvchikov GA. (1986) Alkogolnyi sindrom ploda: (Obzor literatury) [Fetal Alcohol Syndrome: literature review]. *Vopr Okhrany Materinstva i Detstva [Matern Child Prot Issues]* 5:48–54.
- Talykova LV, Vaktskjold A, Serebrjakova NG *et al.* (2007) Pregnancy health and outcome in two cities in the Kola Peninsula, northwestern Russia. *Int J Circumpolar Health* 66:168–81.
- The St. Petersburg-USA Orphanage Research Team. (2005) Characteristics of children, caregivers, and orphanages for young children in St. Petersburg, Russian Federation. *J Appl Dev Psychol* 26:477–506.
- Uliyanovskaia TN, Solovov AG. (2010) Sotsialno-psikhologicheskie osobennosti beremennykh zhenshchin, upotrebliaiushchikh alkogol [Socio-psychological characteristics of pregnant women consuming alcohol]. *Ekologiya Cheloveka [Hum Ecol]* 4:42–5.
- Usova EV, Sazonova NS, Rassadina ZA. (1981) Sindrom alkogolnoi embriopatii [Alcohol embryopathy syndrome]. *Zhurn nevr i Psikiatr im S.S. Korsakova [J Neur Psychiatry named after S.S. Korsakov]* 81:1552–6.
- Vyatchanina ES, Skalny AV. (2009) Izuchenie vliianiia upotrebleniia alkogolia materiu do beremennosti na elementnyi sostav kory golovnogo mozga potomstva (eksperimentalnoe issledovanie) [Research study on the influence of maternal alcohol consumption before pregnancy on the offspring's cortex structure (experimental study)]. V: I Ross. Natsionalnii Kongress po Narkologii s Mezhdunarodnym Uchastiem [In: 1st Rus National Congress on Narcology with International Participation]. Moscow, Russia, 24–27 Nov 2009, pp. 7–8.
- Warren KR, Calhoun FJ, May PA *et al.* (2001) Fetal alcohol syndrome: an international perspective. *Alcohol Clin Exp Res* 25:202S–6S.
- World Health Organization (WHO). (2010) European Status Report on Alcohol and Health 2010. Geneva: WHO Regional Office for Europe. Available at: <http://www.euro.who.int/en/what-we-do/health-topics/disease-prevention/alcohol-use/publications/2010/european-status-report-on-alcohol-and-health-2010> (7 July 2013, date last accessed).
- World Health Organization (WHO). (2011) Global Status Report on Alcohol and Health. Geneva: WHO. Available at: http://www.who.int/substance_abuse/publications/global_alcohol_report/profiles/rus.pdf (7 July 2013, date last accessed).
- Zabludovsky AL, Shikhov NS, Maizelis MY. (1989) Osobennosti povedeniia i uslovnoreflektornoi deiatelnosti zhivotnykh, podvergnutyykh vnutritrubnym vozdeistviyam etanola. V *Posledstviia alkogolnoi intoksikatsii dlia potomstva*, pod red. Airapetiantc MG [Characteristics of behaviour and conditioned reflex activity in animals exposed to ethanol in utero. In Airapetyants MG (ed). *The Consequences of Alcohol Intoxication for Posterity*]. Moskva: Nauka [Moscow: Science], 21–7.
- Zhulin VV, Bazyan AS. (1989) Vliianie vnutritrubnogo vliianiia etanola na nekotorye neiromediatornye sistemy kory mozga krysov. V *Posledstviia alkogolnoi intoksikatsii dlia potomstva*, pod red. Airapetiantc M.G. [Influence of prenatal ethanol exposure on several neurotransmitter systems of the cerebral cortex of rats]. *Posledstviia alkogolnoi intoksikatsii dlia potomstva* [In Airapetyants MG (ed). *The consequences of alcohol intoxication for posterity*]. Moskva: Nauka [Moscow: Science], 15–9.